

## CONCERNING REAL FUNCTIONS WITH VALUES IN THE CANTOR SET

YU-LIN CHOU<sup>1</sup>

**Abstract.** This article in particular indicates that there exist at least continuum-many "essentially nonconstant" (requiring the existence of at least two level sets having positive finite measure) almost everywhere continuous functions from the real field  $\mathbb{R}$  to the Cantor ternary set  $\mathcal{C}$ , although it is a basic fact that there exists no nonconstant continuous function  $\mathbb{R} \rightarrow \mathcal{C}$ .

### 1. INTRODUCTION

A *Cantor space* (i.e. [following [3]], a topological space homeomorphic to the Cantor ternary set), denoted  $\mathcal{C}$  hereafter, topologized as a topological subspace of the real field  $\mathbb{R}$ , seems to be asymmetrically studied as the codomain of a function on  $\mathbb{R}$ . One apparent reason would be the basic fact that every continuous function  $f : \mathbb{R} \rightarrow \mathcal{C}$  is constant: If not, then the  $f$ -image  $f^{-1}(\mathbb{R}) \subset \mathcal{C}$  of  $\mathbb{R}$  has at least two elements; since  $\mathcal{C}$  is Hausdorff and zero-dimensional (i.e. [following [3]], having a basis consisting of clopen sets), it then follows that  $\mathbb{R}$  has a nonempty clopen proper subset, contradicting the fact that  $\mathbb{R}$  is connected.

For our purposes, we introduce the following

**Definition 1.1.** Let  $q \geq 2$  be a positive integer; let  $Y \subset \mathbb{R}$ ; let  $f : \mathbb{R} \rightarrow Y$ . The function  $f$  is said to be essentially ( $q$ -)nonconstant if and only if there exist some distinct  $y_1, \dots, y_q \in Y$  such that the  $f$ -preimage  $f^{-1}(\{y_i\})$  of  $\{y_i\}$  has positive finite (Lebesgue-)measure for all  $1 \leq i \leq q$ .

Thus every essentially nonconstant function is automatically a nonconstant function.

---

2010 *Mathematics Subject Classification.* Primary: 54C20; 26A15.

*Key words and phrases.* almost everywhere continuous extension of continuous map; Borel measure space; Cantor space; dense set; essentially nonconstant map.

In what follows, we give another proof of an "almost everywhere continuous extension" result in a suitably generic setting, which is then applied to show that there are at least continuum-many essentially nonconstant almost everywhere continuous functions  $\mathbb{R} \rightarrow \mathcal{C}$ , in dire contrast to the nonexistence of a nonconstant continuous function  $\mathbb{R} \rightarrow \mathcal{C}$ .

## 2. RESULTS

The continuous extendability of a continuous map from a dense subspace of a topological space to a compact Hausdorff space admits a well-known characterization (e.g., Theorem 3.2.1 in [2]). In view of the inherent restrictions in particular due to this necessary and sufficient condition, such a continuous map need not be continuously extendable in an automatic manner. However, one does have a generic result (Theorem 3.2 in [1]) indicating "how continuous" an extension of such a continuous map can be.

If  $f : X \rightarrow Y$ , we denote by  $f^{\rightarrow}(A)$  the  $f$ -image of  $A$  for all  $A \subset X$ , and by  $f^{\leftarrow}(B)$  the  $f$ -preimage of  $B$  for all  $B \subset Y$ .

We give another proof of Theorem 3.2 in [1] for compact codomains, which is directly relevant with respect to our situation:

**Proposition 2.1.** *Let  $Y$  be a compact Hausdorff space; let  $X$  be a topological space. If  $A \subset X$  is a dense subspace, then every continuous map  $A \rightarrow Y$  is extendable to some map  $g : X \rightarrow Y$  that is continuous at every point of  $A$ .*

*Proof.* Let  $f : A \rightarrow Y$  be continuous. If  $X$  is empty or  $A = X$ , then there is certainly nothing to prove; we consider the other cases. Since  $Y$  is compact, every net in  $Y$  has a cluster point, i.e. (following [2]), for every net  $(y_\theta)_\theta$  in  $Y$  there exists some  $y \in Y$  such that every neighborhood of  $y$  contains  $y_\theta$  frequently in  $\theta$ . On the other hand, by the denseness assumption, for every  $x \in X \setminus A$  we can choose some net  $(a_\theta^x)_{\theta \in \Theta_x}$  in  $A$ , where  $\Theta_x$  is nonempty, converging in  $X$  to  $x$ . Then each net  $(f(a_\theta^x))_{\theta \in \Theta_x}$  in  $Y$  has a cluster point in  $Y$ ; for every  $x \in X \setminus A$ , let  $\widehat{\lim}_{\theta \in \Theta_x} f(a_\theta^x)$  be a cluster point of the net  $(f(a_\theta^x))_{\theta \in \Theta_x}$  by acknowledging the Axiom of Choice.

We claim that the map

$$g : X \rightarrow Y, \begin{cases} x \mapsto f(x), & \text{if } x \in A, \\ x \mapsto \widehat{\lim}_{\theta \in \Theta_x} f(a_\theta^x), & \text{if } x \in X \setminus A \end{cases}$$

is a desired extension of  $f$  over  $X$ . Let  $x \in A$ , and let  $G$  be a neighborhood of  $g(x)$  in  $Y$ . Then, since  $Y$  is by using assumption in particular a regular space, we can choose some neighborhood  $V$  of  $g(x) = f(x)$  such that  $\text{cl}(V) \subset G$ . In turn, we can choose some neighborhood  $W_x$  of  $x$  in  $X$ , by the continuity of  $f$ , such that

$$f^{\rightarrow}(W_x \cap A) \subset V.$$

We show that  $g^{\rightarrow}(W_x) \subset G$ , so that the continuity of  $g$  at  $x$  is verified. To this end, it suffices to show that  $g(x') \in \text{cl}(V)$  for all  $x' \in W_x$ . Let  $x' \in W_x$ . If  $x' \in W_x \cap A$ , then  $g(x') = f(x') \in V$ , and there is nothing to prove. Suppose  $x' \in W_x \cap (X \setminus A)$ , and let  $O$  be a neighborhood of  $g(x')$ . Then the set  $\Theta_{x'}(O) :=$

$\{\theta \in \Theta_{x'} \mid f(a_\theta^{x'}) \in O\}$  is nonempty and cofinal in  $\Theta_{x'}$  by the construction of  $g$ . Since  $(a_\theta^{x'})_{\theta \in \Theta_{x'}}$  converges in  $X$  to  $x'$  by definition, the neighborhood  $W_x$  of  $x'$  contains  $a_\theta^{x'}$  eventually in  $\theta$ . But each  $a_\theta^{x'} \in A$  by definition; it follows that  $f(a_\theta^{x'}) \in V$  eventually in  $\theta$ , and so, in particular, there exists some  $\theta \in \Theta_{x'}(O)$  such that  $f(a_\theta^{x'}) \in V$  and hence  $f(a_\theta^{x'}) \in V \cap O$ . Thus  $g(x') \in \text{cl}(V)$ . We have shown that  $g$  is continuous at every point of  $A$ ; this completes the proof.  $\square$

We record as a relevant corollary of Proposition 2.1 the following evident measure-theoretic application:

**Corollary 0.1.** *Let  $Y$  be a compact Hausdorff space; let  $X$  be a topological space that is also a complete measure space with  $M_X$  denoting the given complete measure and with the given sigma-algebra including the Borel sigma-algebra of  $X$ .*

*If  $A \subset X$  is a dense co-null topological subspace (i.e., if  $A$  is a subset of  $X$ , being dense in the topological space  $X$  and being the complement of some  $M_X$ -null set in the measure space  $X$ , which is topologized as a subspace of the topological space  $X$ ), then every continuous map  $A \rightarrow Y$  can be extended to be some  $M_X$ -almost everywhere continuous map  $X \rightarrow Y$ .*  $\square$

We are now in a position to prove

**Theorem 1.** *For every integer  $q \geq 2$ , there exist at least continuum-many essentially  $q$ -nonconstant almost everywhere continuous functions  $\mathbb{R} \rightarrow \mathcal{C}$ .*

*Proof.* Denote by  $A$  the set of all irrational numbers. Since the subspace  $A$  of  $\mathbb{R}$  is second countable and zero-dimensional, by an elementary argument using the basic fact that being second countable implies being Lindelöf, we can choose some countable basis  $\widehat{\mathcal{T}}_A$  of  $A$  consisting of clopen sets in  $A$ . Upon fixing any  $a_0 \in A$  and choosing a neighborhood  $G$  of  $a_0$  in  $A$  with positive finite measure, e.g.,  $G := ]a_0 - 1, a_0 + 1[ \cap A$ , we can in turn choose some element  $W_1$  of  $\widehat{\mathcal{T}}_A$  such that  $W_1 \subset G$  and  $W_1$  has positive finite measure in view of the countable subadditivity of a measure.

Let  $q \geq 2$  be an integer; we can then choose some distinct  $r_1, \dots, r_{q-1} \in \mathbb{Q}$  such that if  $t_{r_i} : x \mapsto r_i + x$  on  $\mathbb{R}$  for all  $1 \leq i \leq q-1$  then the sets

$$]a_0 - 1, a_0 + 1[, t_{r_1}^\rightarrow(]a_0 - 1, a_0 + 1[), \dots, t_{r_{q-1}}^\rightarrow(]a_0 - 1, a_0 + 1[)$$

are pairwise disjoint. Thus each  $W_{i+1} := t_{r_i}^\rightarrow(W_1)$  has the same positive finite measure as  $W_1$  by the translation invariance of (Lebesgue) measure. Moreover, as each  $t_{r_i}|_A$  is a homeomorphism of  $A$  onto  $A$ , the sets  $W_i$  are all clopen in  $A$ .

If  $c_1 \in \mathcal{C}$ , if  $c_i \in \mathcal{C} \setminus \{c_1, \dots, c_{i-1}\}$  for all  $2 \leq i \leq q$ , and if  $c \in \mathcal{C} \setminus \{c_1, \dots, c_q\}$ , define

$$f : A \rightarrow \mathcal{C}, \begin{cases} a \mapsto c_1, & \text{if } a \in W_1; \\ \vdots & \vdots \\ a \mapsto c_q, & \text{if } a \in W_q; \\ a \mapsto c, & \text{if } a \in A \setminus \bigcup_{i=1}^q W_i. \end{cases}$$

Since the set  $A \setminus \bigcup_{i=1}^q W_i$  is also (cl)open in  $A$ , the map  $f$  is continuous. Then we can choose by Proposition 2.1 some almost everywhere continuous extension  $g : \mathbb{R} \rightarrow \mathcal{C}$  of  $f$ . But  $g$  is essentially  $q$ -nonconstant: We have  $W_1, \dots, W_q$  having positive finite measure by construction, and we have  $c_1, \dots, c_q \in \mathcal{C}$  being distinct; since  $g^{-1}(\{c_i\}) = W_i \cup E$  for some  $E \subset \mathbb{Q}$ , the measure of  $g^{-1}(\{c_i\})$  equals that of  $W_i$  for all  $1 \leq i \leq q$ .

Manifestly, given any finite subset  $Z$  of  $\mathcal{C}$ , the set  $\mathcal{C} \setminus Z$  is in bijection with  $\mathbb{R}$ ; for, we have  $\mathbb{R}$  being equinumerous to  $\mathbb{R} \setminus Z$  and  $\mathbb{R} \setminus Z$  being equinumerous to  $\mathcal{C} \setminus Z$ . Thus the choices of  $f$  as  $c$  runs through  $\mathcal{C} \setminus \{c_1, \dots, c_q\}$ , with each  $c_i$  and each  $W_i$  being fixed, are at least continuum-many; the proof is complete.  $\square$

#### REFERENCES

- [1] C. Costantini, A. Marcone, *Extensions of functions which preserve the continuity on the original domain*, *Topology and its Applications*, 2 (103), Elsevier (2000) 131–153.
- [2] R. Engelking, *General Topology*, revised and completed edition, Heldermann Verlag, Berlin, 1989.
- [3] S. M. Srivastava, *A Course on Borel Sets*, Springer, Berlin, 1998.

HSINCHU COUNTY 310, TAIWAN (R.O.C.)  
 Email address: [chou.y.l.edu@gmail.com](mailto:chou.y.l.edu@gmail.com)

Received 28.9.2023  
 Revised 6.12.2023  
 Accepted 12.12.2023